

**YOUR BOWEL PREP IS EXTREMELY IMPORTANT!  
PLEASE READ INSTRUCTIONS CAREFULLY**

Prep: MoviPrep

<p><b>1 Week Before</b></p>	<ul style="list-style-type: none"> <li>❑ <b>Purchase</b> the following:             <ul style="list-style-type: none"> <li>○ MoviPrep Bowel prep – Pick up from pharmacy.</li> <li>○ GasX or simethicone gas relief over the counter. You will only need 4 of these. These help decrease bubbles in your colon.</li> <li>○ Clear liquids – Refer to clear liquid diet handout.</li> </ul> </li> <li>❑ <b>Plan for a driver:</b> Plan for someone to drive you home or escort you if taking alternate modes of transportation. You will be sedated for your procedure. You should not drive again until the morning after your procedure.</li> <li>❑ <b>Stop eating</b> any visible seeds, nuts or whole grains. You can still eat fruits and vegetables if you remove the seeds from them. Please do not take oil-based vitamins (A, D, E or K, fish oil, etc.) or supplements for a full 7 days before your procedure. While we may not cancel your procedure if you have accidentally eaten them the week before your procedure, it may impact the quality of your prep.</li> <li>❑ If you are <b>diabetic</b> and/or taking <b>prescription blood thinning medication</b>, refer to handout. If you did not receive this handout, contact our office.</li> </ul>
<p><b>1 Day Before</b></p>	<ul style="list-style-type: none"> <li>❑ <b>Start clear liquid diet</b> from the time you wake up. See clear liquid diet handout.</li> <li>❑ <b>In the morning:</b> Empty MoviPrep (one pouch A and one pouch B) into container. Add warm water to the top line and mix well. Place in refrigerator.</li> <li>❑ <b>At 6:00 pm:</b> <ul style="list-style-type: none"> <li>○ Drink MoviPrep solution over the next hour. (8 ounces every 15 minutes).</li> <li>○ <b>Take 2 GasX pills.</b></li> </ul> </li> <li>❑ Continue clear liquids to stay hydrated.</li> <li>❑ <b>In the evening:</b> Empty second MoviPrep (one pouch A and one pouch B) into container. Add warm water to the top line and mix well. Place in refrigerator.</li> </ul>
<p><b>Procedure Day</b></p>	<ul style="list-style-type: none"> <li>❑ <b>Continue clear liquid diet</b> from the time you wake up.</li> <li>❑ <b>6 HOURS PRIOR to your procedure:</b> <ul style="list-style-type: none"> <li>○ Drink second MoviPrep solution over the next hour.</li> <li>○ <b>Take 2 GasX pills.</b></li> <li>○ <b>Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana.</b></li> </ul> </li> <li>❑ <b>5 HOURS PRIOR to your procedure:</b> <ul style="list-style-type: none"> <li>○ Drink at least 2 more 8 ounce glasses of clear liquids over the next hour.</li> </ul> </li> <li>❑ <b>4 HOURS PRIOR to your procedure</b> <ul style="list-style-type: none"> <li>○ NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints.</li> <li>○ The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, <b>BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS).</b></li> </ul> </li> <li>❑ If you cannot see through to the bottom of the toilet, please call our office for further advice after 6 am.</li> <li>❑ <b>Have the following:</b> <ul style="list-style-type: none"> <li>○ A ride home. To be confirmed at check-in.</li> <li>○ Insurance cards and driver's license. Do not bring valuables.</li> </ul> </li> </ul>