

YOUR BOWEL PREP IS EXTREMELY IMPORTANT!

		PLEASE READ INSTRUCTIONS CAREFULLY
Prep: SuTab - 2 Day with	Mad	
Top: Gurus 2 2 ay min		Purchase the following:
	_	O SuTab Bowel prep prescription from your pharmacy.
		• GasX or simethicone gas relief over the counter. You will only need 4 of these.
		These help to reduce bubbles in your GI tract which improves the quality of your
1 Week		exam.
1 VVCCK		O 10-ounce bottle of GREEN or CLEAR magnesium citrate (over the counter laxative).
Before		NO RED OR PURPLE.
Deloie		O Clear liquids – Refer to clear liquid diet handout.
		Plan for a driver: Plan for someone to drive you home or escort you if taking alternate
		modes of transportation. You will be sedated for your procedure. You should not drive
		again until the morning after your procedure.
		Stop eating any visible seeds, nuts or whole grains. You can still eat fruits and
		vegetables if you remove the seeds from them. Refer to the dietary guidelines
		handout. Please do not take oil-based vitamins/supplements (A, D, E or K, fish oil, etc.)
		for a full 7 days before your procedure. While we may not cancel your procedure if you
		have accidentally eaten them the week before your procedure, it may impact the quality
		of your prep.
		If you are diabetic and/or taking prescription blood thinning medication, refer to
		handout. If you did not receive this handout, contact our office.
2 Days		Start clear liquid diet from the time you wake up. Refer to clear liquid diet handout.
2 Days		At 7:00 pm: Drink the 10-ounce bottle of magnesium citrate. Drink over 15 minutes to
Defens		reduce nausea.
Before		Continue clear liquids to stay hydrated.
1 Day Refere		PLEASE FOLLOW THE PREP INSTRUCTIONS PROVIDED BY OUR OFFICE
1 Day Before		Start clear liquid diet from the time you wake up. See clear liquid diet handout.
		At 6:00 pm
		Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink
		the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets.
		the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills.
	_	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it.
	_	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. O If you experience nausea or vomiting take a 15-30 minute break, then continue
		 the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids.
		the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. O If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated.
	<u> </u>	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. O If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up.
		the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. O If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure:
	<u> </u>	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: O Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana.
	<u> </u>	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: O Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. O Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink
	<u> </u>	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. O Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: O Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. O Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes.
	<u> </u>	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills.
Procedure	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour.
_	<u> </u>	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. 4 HOURS PRIOR to your procedure
_	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. 4 HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints.
Procedure Day	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. 4 HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours
_	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. 4 HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP
_	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. 4 HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS).
_	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS). If you cannot see through to the bottom of the toilet one hour after finishing your prep,
_	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS). If you cannot see through to the bottom of the toilet one hour after finishing your prep, please call our office for further advice after 6 am.
_	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS). If you cannot see through to the bottom of the toilet one hour after finishing your prep, please call our office for further advice after 6 am. Have the following:
	0	the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. Take 2 GasX pills. Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. Continue clear liquids to stay hydrated. Continue clear liquid diet from the time you wake up. HOURS PRIOR to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Take 2 GasX pills. Drink two more 16 oz containers of cool/cold water over the next hour. HOURS PRIOR to your procedure NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS). If you cannot see through to the bottom of the toilet one hour after finishing your prep, please call our office for further advice after 6 am.