

SPECIALTY MEDICINE, EXCEPTIONAL CARE

YOUR BOWEL PREP IS EXTREMELY IMPORTANT! PLEASE READ INSTRUCTIONS CAREFULLY

Prep: Clenpiq 3 Day	
1 Week Before	 □ Purchase the following: □ Two Clenpiq Bowel prep – Pick up from pharmacy. □ GasX or simethicone gas relief over the counter. You will only need 4 of these. These help decrease bubbles in your colon. □ Clear liquids – Refer to clear liquid diet handout. □ Plan for a driver: Plan for someone to drive you home or escort you if taking alternate modes of transportation. You will be sedated for your procedure. You should not drive again until the morning after your procedure. □ Stop eating any visible seeds, nuts or whole grains. You can still eat fruits and vegetables if you remove the seeds from them. Refer to the dietary guidelines handout. Please do not take oil-based vitamins (A, D, E or K, fish oil, etc.) or supplements for a full 7 days before your procedure. While we may not cancel your procedure if you have accidentally eaten them the week before your procedure, it may impact the quality of your prep. □ Stop the use of any illicit drugs 7 days prior to your procedure or your procedure may be canceled. □ If you are taking prescription medications, please refer to the Medication Changes handout for holding instructions. If you did not receive this
3 Days Before	handout, contact our office. Start clear liquid diet from the time you wake up. Refer to clear liquid diet handout.
2 Days Before	☐ Continue clear liquid diet from the time you wake up. ☐ At 6:00 pm: ☐ Dripk 1 bettle of CLENDIO prop. pe peed to mix with anything clear
	 Drink 1 bottle of CLENPIQ prep, no need to mix with anything else. After drinking the bottle of prep, drink at least 5 or more 8 oz cups of clear liquid over the next 2 to 4 hours. Take 2 GasX pills.
1 Day Before	Continue clear liquid diet from the time you wake up. At 10:00 am: Drink 1 bottle of CLENPIQ prep, no need to mix with anything else. After drinking the bottle of prep, drink at least 5 or more 8 oz cups of clear liquid over the next 2 to 4 hours
	 At 6:00 pm: Drink 1 bottle of CLENPIQ prep, no need to mix with anything else. After drinking the bottle of prep, drink at least 5 or more 8 oz cups of clear liquid over the next 2 to 4 hours. Take 2 GasX pills.
	 Continue clear liquid diet from the time you wake up. 6 HOURS PRIOR to your procedure: O Drink the LAST bottle of CLENPIQ prep, no need to mix with anything else. O After drinking the bottle of prep, drink at least 5 more 8 oz cups of clear liquid over the next 2 hours.
Procedure	 Take 2 GasX pills. Stop all use of cigarettes, e-cigarettes, chewing tobacco and
Day	marijuana. □ 4 HOURS PRIOR to your procedure:



SPECIALTY MEDICINE, EXCEPTIONAL CARE

 NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints.
O The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS).
If you cannot see through to the bottom of the toilet, please call the number
above for further advice after 7 am.
Have the following:
 A ride home. To be confirmed at check-in.
 Bring Insurance cards and driver's license. Do not bring valuables.