

## SPECIALTY MEDICINE, EXCEPTIONAL CARE

## YOUR BOWEL PREP IS EXTREMELY IMPORTANT! PLEASE READ INSTRUCTIONS CAREFULLY

ried. Sui lave 2 Dav Willi Mauliesiulli Ciliale	Prep: SuFlave 2	Day with	Magnesium	Citrate
---	-----------------	----------	-----------	---------

Prep: Surlave 2 Day with	
	□ Purchase the following:
	SuFlave Bowel Prep- Pick up from your pharmacy. If your insurance does not
	cover this, do not purchase it and contact our office for other options.
1 \\/	O GasX or simethicone gas relief, over the counter. You will only need 4 of
1 Week	these. These help decrease bubbles in your colon.
D - f	o 10-ounce bottle of GREEN or CLEAR magnesium citrate (over the counter
Before	laxative). NO RED OR PURPLE.
	<ul> <li>Clear liquids- Refer to clear liquid diet handout.</li> <li>Plan for a driver: Plan for someone to drive you home or escort you if taking</li> </ul>
	alternate modes of transportation. You will be sedated for your procedure. You
	should not drive again until the morning after your procedure.
	□ Stop eating any visible seeds, nuts or whole grains. You can still eat fruits and
	vegetables if you remove the seeds from them. Refer to the <b>dietary guidelines</b>
	handout. Please do not take any oil-based vitamin/supplements (A, D, E or K, fish
	oil, etc.) for a full 7 days before your procedure. While we may not cancel your
	procedure if you have accidentally consumed them the week before your
	procedure, it may impact the quality of your prep.
	□ Stop the use of any <b>illicit drugs</b> 7 days prior to your procedure or your procedure
	may be canceled.
	☐ If you are taking <b>prescription medications</b> , please refer to the <b>Medication</b>
	Changes handout for holding instructions. If you did not receive this handout,
	contact our office.
2 Days Refere	□ Start clear liquid diet from the time you wake up. Refer to clear liquid diet
2 Days Before	handout.
	□ At 7:00 pm: Drink the 10-ounce bottle of magnesium citrate. Drink over 15 minutes
	to reduce nausea.
	□ Continue clear liquids to stay hydrated.
	Continue clear liquid diet from the time you wake up.
1 Day Before	□ At 5:00 pm:
I Day before	Add one flavor enhancing packet into ONE bottle of SuFlave and fill with
	lukewarm water up to the fill line. Shake the bottle until all powder has
	mixed well (dissolved). Place in refrigerator for 1 hour.
	<ul><li>At 6:00 pm:</li><li>Drink 8 ounces of solution every 15 minutes until the bottle is empty.</li></ul>
	<ul> <li>Drink 8 ounces of solution every 15 minutes until the bottle is empty.</li> <li>Drink an additional 16 oz of water.</li> </ul>
	O Take 2 GasX pills.
	□ Continue clear liquids to stay hydrated.
	□ Continue clear liquid diet from the time you wake up.
	□ 7 HOURS prior to your procedure:
	Add one flavor enhancing packet into ONE bottle of SuFlave and fill with
	lukewarm water up to the fill line. Shake the bottle until all powder has
	mixed well (dissolved). Place in refrigerator for 1 hour.
	□ 6 HOURS prior to your procedure:
Procedure	<ul> <li>Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana.</li> </ul>
Fiocedule	O Drink 8 ounces of solution every 15 minutes until the bottle is empty.
	Brink o danode of delation every to minuted antil the bottle to empty.
Day	O Drink an additional 16 oz of water.
Day	



## SPECIALTY MEDICINE, EXCEPTIONAL CARE

O NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints.	
The only exception: If you have a critical medication that cannot be taken 4-6	
hours early, it can be taken up to 2 hours before the procedure, BUT ONLY	
WITH A SIP OF WATER (no more than 2-3 TABLESPOONS).	
If you cannot see through to the bottom of the toilet one hour after finishing your	
prep, please call the number above for further advice after 7 am.	
□ Have the following:	
A ride home. To be confirmed at check-in.	
<ul> <li>Bring insurance cards and driver's license. Do not bring valuables.</li> </ul>	