

**YOUR BOWEL PREP IS EXTREMELY IMPORTANT!
PLEASE READ INSTRUCTIONS CAREFULLY**

Prep: SuTab 2 Day with Magnesium Citrate

<p>1 Week Before</p>	<ul style="list-style-type: none"> ❑ Purchase the following: <ul style="list-style-type: none"> ○ SuTab Bowel Prep prescription from your pharmacy. ○ GasX or simethicone gas relief over the counter. You will only need 4 of these. These help to reduce bubbles in your GI tract which improves the quality of your exam. ○ 10-ounce bottle of GREEN or CLEAR magnesium citrate (over the counter laxative). NO RED OR PURPLE. ○ Clear liquids- Refer to clear liquid diet handout. ❑ Plan for a driver: Plan for someone to drive you home or escort you if taking alternate modes of transportation. You will be sedated for your procedure. You should not drive again until the morning after your procedure. ❑ Stop eating any visible seeds, nuts or whole grains. You can still eat fruits and vegetables if you remove the seeds from them. Refer to the dietary guidelines handout. Please do not take oil-based vitamins/supplements (A, D, E or K, fish oil, etc.) for a full 7 days before your procedure. While we may not cancel your procedure if you have accidentally consumed them the week before your procedure, it may impact the quality of your prep. ❑ Stop the use of any illicit drugs 7 days prior to your procedure or your procedure may be canceled. ❑ If you are taking prescription medications, please refer to the Medication Changes handout for holding instructions. If you did not receive this handout, contact our office.
<p>2 Days Before</p>	<ul style="list-style-type: none"> ❑ Start clear liquid diet from the time you wake up. Refer to clear liquid diet handout. ❑ At 7:00 pm: Drink the 10-ounce bottle of magnesium citrate. Drink over 15 minutes to reduce nausea. ❑ Continue clear liquids to stay hydrated.
<p>1 Day Before</p>	<p>PLEASE FOLLOW THE PREP INSTRUCTIONS PROVIDED BY OUR OFFICE</p> <ul style="list-style-type: none"> ❑ Start clear liquid diet from the time you wake up. See clear liquid diet handout. ❑ At 6:00 pm <ul style="list-style-type: none"> ○ Open 1 bottle of SuTab tablets and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. Do not cut or crush the tablets. ○ Take 2 GasX pills. ❑ Over the next hour, drink two more 16 oz containers of cool/cold water. Do not guzzle it. <ul style="list-style-type: none"> ○ If you experience nausea or vomiting take a 15-30 minute break, then continue drinking liquids. ❑ Continue clear liquids to stay hydrated.
<p>Procedure</p>	<ul style="list-style-type: none"> ❑ Continue clear liquid diet from the time you wake up. ❑ 6 HOURS prior to your procedure: <ul style="list-style-type: none"> ○ Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. ○ Open 1 bottle of SuTab tablets, and swallow each tablet with a sip of water. Drink the first 16 oz container of water within 20 minutes. ○ Take 2 GasX pills. ○ Drink two more 16 oz containers of cool/cold water over the next hour. ❑ 4 HOURS prior to your procedure: <ul style="list-style-type: none"> ○ NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints.



SPECIALTY MEDICINE, EXCEPTIONAL CARE

Day	<ul style="list-style-type: none">○ The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS).□ If you cannot see through to the bottom of the toilet one hour after finishing your prep, please call the number above for further advice after 7 am.□ Have the following:<ul style="list-style-type: none">○ A ride home. To be confirmed at check-in.○ Bring insurance cards and driver's license. Do not bring valuables.
-----	--