

SPECIALTY MEDICINE, EXCEPTIONAL CARE

YOUR BOWEL PREP IS EXTREMELY IMPORTANT! PLEASE READ INSTRUCTIONS CAREFULLY

Upper Endoscopy	
Prepare for procedure day	 Review the clear liquid diet handout and purchase suggested items as needed. If you did not receive this handout, contact our office. Your clear liquid diet will not begin until 12:00 am (MIDNIGHT) the DAY OF your procedure. Plan for a driver: Plan for someone you know to drive you home or escort you if taking alternate modes of transportation. You will be sedated for your procedure. You should not drive again until the morning after your procedure.
1 Week Before	 Stop the use of any illicit drugs 7 days prior to your procedure or your procedure may be canceled. If you are taking prescription medications, please refer to the Medication Changes handout for holding instructions. If you did not receive this handout, contact our office.
Procedure Day	 At midnight: You may only have clear liquids from now until 4 hours prior to your procedure. You may drink higher protein options such as clear broth or clear protein drinks. No solid foods and NO ALCOHOL. It is important to stay hydrated, so drink plenty of water or other clear liquids. 6 HOURS prior to your procedure: Stop all use of cigarettes, e-cigarettes, chewing tobacco and marijuana. 4 HOURS prior to your procedure: NO DRINKING/NOTHING by mouth – including water, gum, hard candy/mints. Having fluids in your stomach, even a lot of saliva, puts you at risk for vomiting and possibly inhaling fluids into your lungs while you are sedated. The only exception: If you have a critical medication that cannot be taken 4-6 hours early, it can be taken up to 2 hours before the procedure, BUT ONLY WITH A SIP OF WATER (no more than 2-3 TABLESPOONS). For your safety, failure to follow this will result in your procedure being cancelled and rescheduled. NO EXCEPTIONS. Have the following: